



MULTIVITAMINS

Per tablet contains:

Thiamine Mononitrate (Vitamin B ₁).....	50 mg
Riboflavin (Vitamin B ₂).....	20 mg
Pyridoxine Hydrochloride (Vitamin B ₆).....	5 mg
Cyanocobalamin (Vitamin B ₁₂).....	5 mcg
Nicotinamide.....	50 mg
Calcium Pantothenate.....	20 mg
Ascorbic Acid (Vitamin C).....	500 mg

ENERVON[®] TABLET

PRODUCT DESCRIPTION

This product is an orange-yellow, capsule-shaped, sugar-coated tablet with a glossy surface.

WHAT IS IN THE MEDICINE?

This product is especially formulated for adults to help ensure optimum energy and increase body resistance against infections. It contains B-complex vitamins (Vitamins B₁, B₂, B₆, B₁₂, Nicotinamide, and Calcium Pantothenate) to help optimize conversion of food into energy that the body can utilize for numerous physiologic processes such as respiration, digestion, blood circulation, immune system response, and as fuel for physical activities. It also has Vitamin C, an antioxidant which helps the body's natural defense against free radicals and help boost immune function.

STRENGTH OF THE MEDICINE

Please see formulation

WHAT IS THE MEDICINE USED FOR?

- A nutritional supplement to help promote increased energy and enhance the immune system
- For the treatment of Vitamin B-complex deficiencies and Vitamin C deficiencies.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THE MEDICINE?

Orally, one tablet daily.

Or, as directed by a doctor.

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

If you are allergic to any ingredient of the product.

UNDESIRABLE EFFECTS

Nicotinamide

- Dosages of Nicotinamide in excess of 100 mg daily may cause flushing of the face, arms, and chest, itching, and nausea.

Vitamin C

- Prolonged intake of Ascorbic Acid (Vitamin C) in excess of 2 grams per day may lead to nausea, abdominal cramps, diarrhea, and nose bleeds.

WHAT OTHER MEDICINE OR FOODS SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

No clinically significant interaction is expected with the amounts of nutrients in the formulation when used as directed.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose, just give the next dose and the subsequent doses at the usual recommended schedule.

Do not double the dose.

HOW SHOULD YOU KEEP THE MEDICINE?

- Keep the product out of reach and sight of children.
- Store at temperatures not exceeding 30°C.

SIGNS AND SYMPTOMS OF OVERDOSAGE

The B-complex Vitamins (Vitamins B₁, B₂, B₆, B₁₂, Nicotinamide, and Calcium Pantothenate) and Vitamin C are relatively nontoxic.

However, long term (i.e., two months or longer) administration of large (megadose) dosages (e.g., usually 2 grams or more daily) of Vitamin B₆ can cause neurological symptoms manifested as paresthesia (more noticeable at night and limited to the extremities), bone pains, hyperesthesia (described as burning, pricking, stinging, or itching), muscle weakness, fasciculation (described as twitching, restlessness or fidgeting), and numbness on the limbs and face.

WHAT TO DO WHEN YOU HAVE USED MORE THAN THE RECOMMENDED DOSAGE?

If you have taken more than the recommended dosage, consult a doctor or contact a Poison Control Center immediately.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- Do not take more than the recommended dose.
- Do not use after the expiry date on the label.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

If any undesirable effect occurs.

AVAILABILITY

Bottle of 45's & 30's tablets, and box of 100's & 8's tablets (in flex foil x 4's)

Manufactured by United Laboratories, Inc.
66 United St., Mandaluyong City, Philippines
for UNITED AMERICAN PHARMACEUTICALS, INC.
132 Pioneer Street, Mandaluyong City, Philippines



Trusted Quality Healthcare

P30000010771

Reg. IPOPHIL